

Dissertação - Artigo de investigação médica

Mestrado Integrado em Medicina – ICBAS-UP

Effects of gradual change of Exercise Intensity on Muscle Oxygenation, measured by Near-infrared Spectroscopy

Autor: Nuno Miguel Machado Amorim

Orientador: Dr. Pedro de Pinho e Costa Amorim, HGSA-CHP

Co-orientador: Prof. Doutor Álvaro José Barbosa Moreira da Silva, HGSA-CHP

Acknowledgments

Em primeiro lugar, agradeço a todos os voluntários que participaram no estudo, que demonstraram sempre muita vontade em ajudar a investigação e curiosidade em saber os resultados deste estudo inovador.

Em segundo lugar, quero agradecer ao Dr. Pedro Amorim por me ter dado a oportunidade de trabalhar com ele durante o último ano. Obrigado pelo voto de confiança. Lembro-me do dia em que me fez a sugestão do tema: fiquei rendido à sua inovação no mundo do desporto. Agora, sinto-me ainda mais entusiasmado por saber que o trabalho terá continuidade.

Em terceiro lugar, quero agradecer ao Dr. Fernando Correia por estar na base do projecto com o Dr. Pedro Amorim e se preocupar constantemente com a evolução do estudo. Obrigado pelas maratonas de tratamento de dados, principalmente a “Maratona Casa Guedes”. Esteve sempre disponível, o que me ajudou bastante a perder o nervosismo na parte final, nas datas de entrega da Tese.

Quero também agradecer ao Dr. Jaime Milheiro e à CMEP, pela disponibilidade, simpatia e entusiasmo que demonstraram em investigar connosco. O Dr. Jaime contribui de uma forma essencial para o estudo, tendo estado presente em todas as avaliações e conduzido todos os testes que foram realizados.

Agradeço à Medtronic pela cedência do aparelho INVOS e dos sensores de NIRS. Foi um contributo essencial para a realização do estudo.

Um agradecimento muito especial aos meus pais e à minha irmã por serem o meu suporte diário. Tudo o que sou agora e todo o meu sucesso deve-se a vocês. E, por essa razão, isto é também vosso.

Muito obrigado, Sofia. Obrigado por seres a melhor do mundo! Um apoio incansável. Apesar da distância física que nos separa, sinto que estás sempre comigo e isso é sem dúvida o mais importante. Quem diria há cinco anos que celebraríamos esta etapa juntos...

Quero agradecer aos meus avós pelo amor e carinho que me transmitem em todos os momentos que estamos juntos. Vocês são um exemplo de união que sigo. Sei que estão a ver um médico a formar-se pela primeira vez na família e espero conseguir encher-vos de orgulho todos os dias.

Por último, quero agradecer aos meus amigos. São também vocês peças importantíssimas neste percurso, principalmente por me fazerem rir todos os dias. “Os amigos são a família que eu escolhi.”

Resumo

Introdução

A intensidade do exercício em ciclistas, assim como a variação da cadência são factores muito importantes na oxigenação muscular dos mesmos. O objectivo do estudo foi avaliar a relação do aumento gradual da intensidade do exercício de pedalagem na oxigenação muscular em ciclistas bem como a fase recuperação após exaustão, comparando um grupo de profissionais e amadores, utilizando a Espectroscopia de Infravermelho próximo (NIRS).

Métodos

Dois grupos de atletas caucasianos do sexo masculino (Grupo A: três atletas profissionais; Grupo B: três atletas amadores) pedalarão até à exaustão, realizando um protocolo incremental, usando o Ergómetro de laboratório Lode Excalibur Sport e os próprios pedais. A cadência permaneceu constante e a potência foi aumentando ao longo do exercício. A oxigenação muscular foi monitorizada continuamente pela Espectroscopia de Infravermelho próximo (NIRS) no músculo Vasto Lateral (VL), e os parâmetros hemodinâmicos (Ventilação [VE], *Uptake* O₂ [VO₂], *Output* de CO₂ [VCO₂] e Frequência Cardíaca [HR]) foram monitorizados continuamente por um aparelho automático de análise e electrocardiograma.

Resultados

Durante o exercício, quando a intensidade aumenta, o músculo diminui a sua saturação de O₂ de uma forma linear, em atletas profissionais e amadores, assim como a Frequência Cardíaca e o VO₂/kg. Quanto à Ventilação, há um aumento quadrático em relação à intensidade, em ambos os grupos. Durante a recuperação, a saturação de O₂ aumenta de uma forma linear.

Conclusões

Durante o teste, a saturação de O₂ muscular descreve uma descida linear, em ambos os grupos.. A velocidade de recuperação da oxigenação muscular após a exaustão é significativamente superior nos ciclistas profissionais, quando comparada com os ciclistas amadores. Por fim, isto leva-nos a concluir que uma maior variação relativa da extracção muscular de O₂ durante o exercício em atletas profissionais, sugere uma produção aumentada de energia aeróbia, e consequentemente maior aptidão aeróbia. Estes são, então, factores muito importantes na saturação muscular de O₂.

Palavras-chave

Oxigenação muscular, NIRS, ciclismo, Saturação O₂.

Abstract

Background

Exercise intensity in cycling, as well cadence variation, is one of the most important factors in cyclists muscle oxygenation. The aim of the study was to assess the relationship between pedaling exercise intensity and muscle oxygenation in cyclists as well as the recovery phase after exhaustion, measured by Near-Infrared Spectroscopy (NIRS), comparing a group of professional cyclists and a group of amateur cyclists.

Methods

Two groups of male caucasian athletes (Group A: three professional cyclists; Group B: three amateur cyclists) cycled until exhaustion, performing an incremental protocol, using the Lode Excalibur Sport Ergometer laboratory bicycle and their own pedals. The pedaling cadence remained constant and the working power changed throughout the exercise. Muscle oxygenation was monitored continuously by near-infrared spectroscopy on the *Vastus Lateralis*, and Hemodynamic parameters (Ventilation [VE], O₂ uptake [VO₂], CO₂ Output [VCO₂] e Heart Rate [HR]) were monitored continuously along the test by an automatic analyzer and Electrocardiogram.

Results

During incremental exercise, muscle O₂ saturation describes a linear decrease, in both groups. HR and VO₂/kg increase in a linear way during exercise, but VE increases in a quadratic way in professional and amateur cyclists. During muscle saturation recovery, the linear increase of O₂ saturation is significantly different between the two groups.

Conclusions

During exercise, muscle O₂ saturation has a linear decrease, in both groups. The rate of muscle O₂ saturation recovery after exhaustion is significantly higher in professional cyclists, when compared to amateur cyclists. This last finding and the larger relative change of muscle O₂ extraction in the working muscle throughout the test in professional cyclists, suggest that higher aerobic energy production, and consequently higher aerobic fitness. These are very important factors in muscle O₂ saturation.

Keywords

Muscle oxygenation, NIRS, cycling, O₂ saturation.

Introduction

The impacts of the changes in pedaling cadence on performance, cardiopulmonary and metabolic responses have attracted the attention of researchers and it is known that time until exhaustion decreases with the increase of the pedaling cadence [1-5], especially in untrained subjects, but not in triathletes. The key link between pedaling cadence and performance seems to be the aerobic fitness of the subject [6]. A decrease in muscle efficiency at high cadences has been described as the main reason for those observations [2, 4, 5, 7, 8]. Concerning the pulmonary oxygen uptake, contradictory results were obtained: while some studies showed no relationship between pedal cadence and pulmonary oxygen uptake [3, 9-11] others claimed the occurrence of higher pulmonary oxygen uptake at higher pedal cadence for the same work rate [1, 2, 4, 8, 12, 13]. Accordingly, the reason why professional cyclists spontaneously adopt higher cadences (over 90 rpm) during endurance cycling, flat stages and time trials is not fully explained by energy consumption. It is also reported that the most economical cadence increases with increasing workload in elite cyclists [10, 14], which is supported by a lower peak force and shorter duration for muscle contraction, which may cause a reduction in intramuscular pressure, alleviating blood flow restriction and preventing muscle fatigue [15, 16]. Literature has accepted these reasons to justify the higher cadences adopted spontaneously by professional cyclists during endurance cycling, flat stages and time trials [17]. Also, a greater glycogen depletion in type II fibers at 50 rpm, compared to 100 rpm, at high workloads has been described [18]. This indicates a greater activation of the quadriceps muscle at 50 rpm in opposition to 100 rpm, explaining the possible benefit of choosing higher cadences at high workloads. The finding that minimum EMG amplitude occurs at a progressively higher cadence as power output supports this theory [19].

The large monoarticular muscles working over the hip and knee joints are considered the main power producing muscles (the gluteus and the vasti muscles) [20]. The *Vastus Lateralis* (VL), among the vasti muscles, is one of the main contributors to pedaling work force in pedaling exercise, so it is commonly used for NIRS studies during pedaling exercise [17, 21].

Cycling research has been focused on studying the effects of changing pedaling cadence on performance, pulmonary oxygen uptake or muscle oxygenation, while keeping exercise power constant. To our knowledge, no study had observed the relationship between muscle oxygenation and the change of working power, during an incremental exercise until exhaustion. We designed the present study to assess the effects of a gradual change of exercise intensity on muscle oxygenation, in professional and amateur cyclists, with the objective of obtaining new

knowledge that could be implemented in a protocol using non-invasive NIRS technology to individualize athletes' training methodologies.

Methods

Subjects

Two groups of subjects were studied: a group of three male professional cyclists with a 2016 activity of ≥ 20000 km (age $28,7 \pm 6,4$ years, weight $67,3 \pm 4,5$ kg, height $177 \pm 3,6$ cm, $20,7 \pm 0,6$ kg/m² BMI); and a group of three male amateur cyclists (age $34 \pm 3,6$ years, weight $71,7 \pm 6,1$ kg, height $181 \pm 3,4$ cm, $22 \pm 2,6$ kg/m² BMI) with a 2016 activity between 5000-10000 km and last month's activity ≥ 400 km. None of the subjects reported a history of injury during the previous 2 years. Before the test, all subjects received a detailed explanation of the experimental protocol, they were informed about the risks and signed the statement of consent. The study was approved by the Clínica Médica do Exercício do Porto (CMEP) Ethics Department.

Experimental protocol

Each subject from the first group performed an incremental cycle exercise test until exhaustion (RAMPA 45) to determine VO₂ max and the power output achieved at VO₂ max. The test began with 3 minutes resting, followed by 3 minutes of warming at 0 W. After that, the power was increased constantly with a rate of 45 W per minute until exhaustion. Pedal cadence was kept constant between 80-90 rpm and the test was stopped manually when the subject was not able to keep increasing the VO₂, despite strong verbal encouragement. After that, the test finished with 3 min of active recovery, cycling at 0 W. The second group performed the same incremental exercise test until exhaustion, but with a power increase rate of 35 W per minute (RAMPA 35). The Lode Excalibur Sport Ergometer (The Netherlands), which is renowned worldwide as the *gold-standard* in cycling ergometry, was used to this experiment, to measure the cycling power.

Measurements

Hemodynamic parameters

An automatic analyzer (Metalyzer 3B, Cortex, Germany) was used to assess the gas exchange variables during each test, and a facial mask with gas detector sensor connected to the analyzer was applied to each participant. It was calibrated according to manufacturer's recommendations with gases of known concentration (16% O₂ and 4% CO₂) before the tests. A 12-lead electrocardiogram was used to monitor continuously the heart rate (HR). Pulmonary oxygen uptake (VO₂), Carbon dioxide output (VCO₂), ventilation (VE) and HR values were averaged into 15s intervals. Data was automatically displayed in the software MetaSoft Studio 3.0 (Cortex, Germany). Anaerobic Threshold (AT) is the moment when VCO₂ changes the increasing rate, when the slope of VCO₂ increases, was manually determined.

Muscle oxygenation

Local muscle oxygenation was measured continuously with a continuous-wave NIRS system (Invos Cerebral/Somatic Oximeter, Medtronic, USA) in the VL during the test. The NIRS probe is composed of two light sources and one photodiode detector, which were spaced by 3 cm and 4 cm, to provide a light penetration depth of 2 cm [22], which allows to have an actual measure of oxygenation at the level of VL. The sensor was applied to the shaved skin, overlying the VL, 10 cm above the upper border of the patella, and parallel to the major axis of the thigh. It was fixed with tape to avoid movement during the test and covered with the cycling shorts, to avoid contamination of external light to the measurement photon pathway. Before each test, the device was calibrated according to the manufacturer's recommendations. The large monoarticular muscles working over the hip and knee joints are considered the main power producing muscles (the gluteus and the vasti muscles) [20]. The Vastus Lateralis, among the vasti muscles, is one of the main contributors to pedaling work force in pedaling exercise, so it is commonly used for NIRS studies during pedaling exercise [17, 21]. NIR light (700-1000 nm) penetrates superficial layers (skin, subcutaneous fat, skull, etc.) and is either absorbed by chromophores (oxy- and deoxyhemoglobin and myoglobin) or scattered within the tissue. NIRS is a noninvasive and relatively low-cost optical technique that is becoming a widely used instrument for measuring tissue O₂ saturation, changes in hemoglobin volume and, indirectly, brain/muscle blood flow and muscle O₂ consumption. This allows us to investigate the balance between O₂ delivery and extraction at the muscle level [23, 24].

The NIRS device provides the mean of the absolute values for oxygenation level in each 6 seconds of the test. Consequently, we decided to express and analyze the relative values against the higher value during the warming phase, for each participant, which we considered as 100%.

Statistics

The difference between two independent experimental groups was evaluated using the unpaired student t-test for normally distributed variables. The significance level was set for all statistical analyses at $p < 0,05$. Data analysis was performed with the GraphPad Prism 7 software for Mac.

Results

Hemodynamic parameters

HR increased over time and power increasing in all volunteers. During the recovery time, HR decreased in both groups. As shown in Figure 1, inter-group comparison revealed that HR is significantly different during exercise until exhaustion ($p < 0,001$). HR increases with power in a linear way (Group A: $r^2 = 0,93$; Group B: $r^2 = 0,84$).




Figure 1 – Relationship between Heart rate in both groups (HR2) and Power (Comparison between the two groups)

On the other hand, VE showed an increase in a quadratic way in both groups (Group A: $r^2=0,92$, $p<0,0001$; Group B: $r^2=0,97$, $p<0,0001$). The slopes of each regression are significantly different ($p<0,0001$). (Figure 2)




Figure 2- Relationship between VE and Power (Comparison between the two groups)

VO_2/kg was calculated and we found that it increases during exercise (Group A: $r^2= 0,91$; Group B: $r^2=0,97$). VO_2/kg along the exercise is significantly different between the two groups ($p=0,0183$). (Figure 3)




Figure 3 – Relationship between VO_2/kg and Power (Comparison between the two groups)

Muscle oxygenation

	Pro cyclists	Amateur cyclists
Rest	73,0	88,2
Start pedaling	86,6	89,8
Start power	95,1	89,7
AT	50,8	76,0
Final Load	10,9	55,4

Table I – Relative Muscle O₂ Saturation in distinct phases – Mean values (%).

As shown in Table I, professional cyclists have a greater relative deoxygenation during exercise, when both groups are compared. At the AT point, professionals reveal a lower muscle O₂ saturation, as well as in the final load.

All the participants leveled their final load intensity at 10 in The Borg Scale [25] which is a simple method of rating perceived exertion.

Muscle O₂ saturation decreased during the incremental exercise and increased abruptly during the recovery, in all the subjects from both groups. (Figure 4)

Values of O₂ saturation are relative to the highest O₂ saturation during warming phase.




Figure 4 – Relative Muscle O₂ saturation over time.

As shown in Figure 5, all the regressions have a r^2 between 0,88 and 0,96. Intra-group comparison showed different slopes of desaturation but the mean slopes from each group were not significantly different ($p=0,21$) (Figure 6).




Figure 5 – Relationship between Relative Muscle O₂ Saturation during incremental exercise.




Figure 6 – Muscle Desaturation slopes in Pro and Amateur cyclists.

On the other hand, inter-group comparison showed that the muscle saturation recovery is significantly faster in professional athletes compared to amateur athletes ($p=0,047$) (Figure 7).




Figure 7 – Relative Muscle O2 Saturation Recovery.




Figure 8 – Muscle Desaturation slopes in Pro and Amateur cyclists.

Discussion

To our knowledge, this is the first study that compares both gradual increment of working power and the effect of aerobic fitness (professional vs. amateur cyclists) on the response of muscle oxygenation during exercise until exhaustion and recovery phase. The main findings of this study were (1) during incremental exercise performed until exhaustion, muscle oxygenation shows a linear decrease in professional and amateur cyclists, and (2) during the recovery phase, the muscle O₂ saturation is significantly different between groups, more specifically, the muscle O₂ saturation recovery rate is significantly higher in professional cyclists, when compared to amateur cyclists. This finding supports the idea of an important contribution of aerobic fitness and muscle condition in performance [6].

We observed a higher acceleration of VO₂/kg in professional athletes, comparing to amateur athletes, during the incremental test. However, the HR has the opposite behavior, with higher increasing rate in amateurs. In addition, concerning the VE, amateurs also have a higher increasing rate, when compared to professionals, which suggests lower aerobic fitness.

We observed that professional cyclists have a larger relative change of muscle O₂ extraction in the working muscle throughout the test, when compared with amateur cyclists (Table I).

Professional cyclists, who have higher cycling experience, do not experience an increase of anaerobic energy production during the decrease in the oxygen level during the test [26], demonstrated by NIRS values. In fact, what happens is the dissociation of O₂ from hemoglobin and myoglobin, regarding aerobic metabolism in mitochondria in the VL. It is already explained the effects of endurance training on the increasing of muscle oxidative capacity [27-29] by means of the increase in capillary density (enhanced muscle perfusion), in slow twitch fiber area, and enzyme activity [28, 30, 31]. We do not have any biochemical or histological analyses from our participants, however, the participants' aerobic power, cycling experience and competitive performance suggest that professional athletes have already acquired such developments, contributing to an increase of muscle oxidative capacity through their daily training routine.

Therefore, the capacity is higher in professional than in amateur cyclists. Additionally, the pedaling skill also influences the muscle physiology, because higher skill reduces stress on the leg extensor muscle while cycling, even in highly trained cyclists [32]. These observations can explain, in part, the higher rate in muscle O₂ saturation recovery in professional cyclists: during this phase, the muscle alleviates the pressure instantly and reduces the O₂ extraction and the developments mentioned above cause a rapidly recovering of the baseline muscle saturation.

NIRS technology has been used in sports medicine, namely in dynamic exercise. These findings may allow the study and the improvement of training methodologies and its efficiency.

Methodological considerations

The main limitation of the study is the lack of sensitivity of the NIRS system to muscle O₂ saturations below 15%, during heavy exercise. The system is not designed to measure such low saturations.

The other limitation is that the study results concern only a small part of the VL (4 cm), and can not be assumed to other muscles involved in cycling. However, from all the muscles involved, VL is the one that gives a better representation, making these results very interesting.

Prior the beginning of one of the tests, the gas analyzer did not calibrate completely. Data from that subject gave us values a little higher than expected.

Conclusions

In summary, during incremental exercise performed until exhaustion, muscle deoxygenation decreased in a linear way in both groups. Professional cyclists had a higher VO₂ max, reached a higher heart rate and pedaled until a higher load, as expected. They did so at significantly lower muscle oxygenation levels. Amateur cyclists had much higher muscle oxygenation, but in spite of that they could not achieve high exercise loads. This suggests that the ability of achieving very low levels of muscle oxygenation is an indicator of aerobic fitness and not the opposite. Reaching very low muscle oxygenation levels with exercise does not imply lack of fitness. Professional cyclists can perform at anaerobic levels. Muscle oxygenation recovers faster in athletes. Still it is puzzling that at rest athletes have lower muscle oxygen values. Second, during the recovery phase, the muscle O₂ saturation is significantly different between groups, more specifically, the muscle O₂ saturation recovery rate is significantly lower in amateur cyclists, when compared to professional cyclists. This last finding and the larger relative change of muscle O₂ extraction in the working muscle throughout the test in professional cyclists, suggest that higher aerobic energy production, and consequently higher aerobic fitness.

References

1. Coast, J.R. and H.G. Welch, *Linear increase in optimal pedal rate with increased power output in cycle ergometry*. Eur J Appl Physiol Occup Physiol, 1985. **53**(4): p. 339-42.
2. Gaesser, G.A. and G.A. Brooks, *Muscular efficiency during steady-rate exercise: effects of speed and work rate*. J Appl Physiol, 1975. **38**(6): p. 1132-9.
3. Hill, D.W. and J.L. Vingren, *The effect of pedalling cadence on maximal accumulated oxygen deficit*. Eur J Appl Physiol, 2012. **112**(7): p. 2637-43.
4. Nesi, X., et al., *Effect of a 15% increase in preferred pedal rate on time to exhaustion during heavy exercise*. Can J Appl Physiol, 2004. **29**(2): p. 146-56.
5. Nielsen, J.S., E.A. Hansen, and G. Sjogaard, *Pedalling rate affects endurance performance during high-intensity cycling*. Eur J Appl Physiol, 2004. **92**(1-2): p. 114-20.
6. Zorgati, H., et al., *Effect of pedaling cadence on muscle oxygenation during high-intensity cycling until exhaustion: a comparison between untrained subjects and triathletes*. Eur J Appl Physiol, 2015. **115**(12): p. 2681-9.
7. Sargeant, A.J., *Human power output and muscle fatigue*. Int J Sports Med, 1994. **15**(3): p. 116-21.
8. Tokui, M. and K. Hirakoba, *Estimation of oxygen cost of internal power during cycling exercise with changing pedal rate*. J Physiol Anthropol, 2008. **27**(3): p. 133-8.
9. Barstow, T.J., et al., *Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise*. J Appl Physiol (1985), 1996. **81**(4): p. 1642-50.
10. Foss, O. and J. Hallen, *The most economical cadence increases with increasing workload*. Eur J Appl Physiol, 2004. **92**(4-5): p. 443-51.
11. Zoladz, J.A., et al., *Effect of different cycling frequencies during incremental exercise on the venous plasma potassium concentration in humans*. Physiol Res, 2002. **51**(6): p. 581-6.
12. Chavarren, J. and J.A. Calbet, *Cycling efficiency and pedalling frequency in road cyclists*. Eur J Appl Physiol Occup Physiol, 1999. **80**(6): p. 555-63.
13. Zorgati, H., et al., *Effect of pedal cadence on the heterogeneity of muscle deoxygenation during moderate exercise*. Appl Physiol Nutr Metab, 2013. **38**(12): p. 1206-10.
14. Boone, J., et al., *The impact of pedal rate on muscle oxygenation, muscle activation and whole-body VO(2) during ramp exercise in healthy subjects*. Eur J Appl Physiol, 2015. **115**(1): p. 57-70.
15. Takaishi, T., et al., *Neuromuscular, metabolic, and kinetic adaptations for skilled pedaling performance in cyclists*. Med Sci Sports Exerc, 1998. **30**(3): p. 442-9.
16. Lollgen, H., T. Graham, and G. Sjogaard, *Muscle metabolites, force, and perceived exertion bicycling at varying pedal rates*. Med Sci Sports Exerc, 1980. **12**(5): p. 345-51.
17. Lucia, A., J. Hoyos, and J.L. Chicharro, *Preferred pedalling cadence in professional cycling*. Med Sci Sports Exerc, 2001. **33**(8): p. 1361-6.
18. Ahlquist, L.E., et al., *The effect of pedaling frequency on glycogen depletion rates in type I and type II quadriceps muscle fibers during submaximal cycling exercise*. Eur J Appl Physiol Occup Physiol, 1992. **65**(4): p. 360-4.
19. MacIntosh, B.R., R.R. Neptune, and J.F. Horton, *Cadence, power, and muscle activation in cycle ergometry*. Med Sci Sports Exerc, 2000. **32**(7): p. 1281-7.
20. Skovereng, K., G. Ettema, and M.C.P. van Beekvelt, *Oxygenation, local muscle oxygen consumption and joint specific power in cycling: the effect of cadence at a constant external work rate*. European Journal of Applied Physiology, 2016. **116**: p. 1207-1217.
21. Wang, B., et al., *Differences between the Vastus Lateralis and Gastrocnemius Lateralis in the Assessment Ability of Breakpoints of Muscle Oxygenation for Aerobic Capacity Indices During an Incremental Cycling Exercise*. Journal of Sports Science & Medicine, 2012. **11**(4): p. 606-613.

22. Bhambhani, Y., R. Malik, and S. Mookerjee, *Cerebral oxygenation declines at exercise intensities above the respiratory compensation threshold*. Respir Physiol Neurobiol, 2007. **156**(2): p. 196-202.
23. Ferrari, M., L. Mottola, and V. Quaresima, *Principles, techniques, and limitations of near infrared spectroscopy*. Can J Appl Physiol, 2004. **29**(4): p. 463-87.
24. Scheeren, T.W., P. Schober, and L.A. Schwarte, *Monitoring tissue oxygenation by near infrared spectroscopy (NIRS): background and current applications*. J Clin Monit Comput, 2012. **26**(4): p. 279-87.
25. Borg, G.A., *Psychophysical bases of perceived exertion*. Med Sci Sports Exerc, 1982. **14**(5): p. 377-81.
26. Takaishi, T., et al., *Effect of cycling experience and pedal cadence on the near-infrared spectroscopy parameters*. Med Sci Sports Exerc, 2002. **34**(12): p. 2062-71.
27. Holloszy, J.O. and E.F. Coyle, *Adaptations of skeletal muscle to endurance exercise and their metabolic consequences*. J Appl Physiol Respir Environ Exerc Physiol, 1984. **56**(4): p. 831-8.
28. Henriksson, J., *Training induced adaptation of skeletal muscle and metabolism during submaximal exercise*. The Journal of Physiology, 1977. **270**(3): p. 661-675.
29. Saltin, B., et al., *The nature of the training response; peripheral and central adaptations of one-legged exercise*. Acta Physiol Scand, 1976. **96**(3): p. 289-305.
30. Brodal, P., F. Ingjer, and L. Hermansen, *Capillary supply of skeletal muscle fibers in untrained and endurance-trained men*. Am J Physiol, 1977. **232**(6): p. H705-12.
31. Klausen, K., L.B. Andersen, and I. Pelle, *Adaptive changes in work capacity, skeletal muscle capillarization and enzyme levels during training and detraining*. Acta Physiol Scand, 1981. **113**(1): p. 9-16.
32. Coyle, E.F., et al., *Determinants of endurance in well-trained cyclists*. J Appl Physiol (1985), 1988. **64**(6): p. 2622-30.